

TAME AUTOMORPHISMS OF $\mathbb{C}^3$ WITH MULTIDEGREE OF THE FORM (p_1, p_2, d_3)

MAREK KARASÍ

ABSTRACT. Let $d_3 \geq p_2 > p_1 \geq 3$ be integers such that p_1, p_2 are prime numbers. In this paper we show that the sequence (p_1, p_2, d_3) is the multidegree of some tame automorphisms of $\mathbb{C}^3$ if and only if $d_3 \in p_1\mathbb{N} + p_2\mathbb{N}$, i.e. if and only if d_3 is a linear combination of p_1 and p_2 with coefficients in $\mathbb{N}$.

1. INTRODUCTION

Let $F = (F_1, \dots, F_n) : \mathbb{C}^n \rightarrow \mathbb{C}^n$ be any polynomial mapping. By multidegree, denoted $\text{mdeg } F$, we call the sequence of positive integers $(\deg F_1, \dots, \deg F_n)$. In dimension 2 there is a complete characterization of the sequences (d_1, d_2) such that there is a polynomial automorphism $F : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ with $\text{mdeg } F = (d_1, d_2)$. This characterization is a consequence of the Jung van der Kulk theorem [1, 4]. Moreover in [3] it was proven, among other things, that there is no tame automorphism of $\mathbb{C}^3$ with multidegree $(3, 4, 5)$, $(3, 5, 7)$, $(4, 5, 7)$ and $(4, 5, 11)$.

Recall that a tame automorphism is, by definition, a composition of linear automorphisms and triangular automorphisms, where a triangular automorphism is a mapping of the following form

$$T : \mathbb{C}^n \ni \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \mapsto \begin{pmatrix} x_1 \\ x_2 + f_2(x_1) \\ \vdots \\ x_n + f_n(x_1, \dots, x_{n-1}) \end{pmatrix} \in \mathbb{C}^n.$$

By $\text{Tame}(\mathbb{C}^n)$ we will denote the group of all tame automorphisms of $\mathbb{C}^n$, and by mdeg the mapping from the set of all polynomial endomorphisms of $\mathbb{C}^n$ into the set $\mathbb{N}^n$. Using this notation, above mentioned facts can be written as follows $(3, 4, 5), (3, 5, 7), (4, 5, 7), (4, 5, 11) \notin \text{mdeg}(\text{Tame}(\mathbb{C}^3))$.

In this paper we make a next progress in the investigation of the set $\text{mdeg}(\text{Tame}(\mathbb{C}^3))$. Namely we show the following theorem.

Theorem 1. *Let $d_3 \geq p_2 > p_1 \geq 3$ be positive integers. If p_1 and p_2 are prime numbers, then $(p_1, p_2, d_3) \in \text{mdeg}(\text{Tame}(\mathbb{C}^3))$ if and only if $d_3 \in p_1\mathbb{N} + p_2\mathbb{N}$, i.e. if and only if d_3 is a linear combination of p_1 and p_2 with coefficients in $\mathbb{N}$.*

Notice that for all permutation σ of the set $\{1, 2, 3\}$, $(d_1, d_2, d_3) \in \text{mdeg}(\text{Tame}(\mathbb{C}^3))$ if and only if $(d_{\sigma(1)}, d_{\sigma(2)}, d_{\sigma(3)}) \in \text{mdeg}(\text{Tame}(\mathbb{C}^3))$. Since, also, $(d_1, d_2, d_3) \in \text{mdeg}(\text{Tame}(\mathbb{C}^3))$ if $d_1 = d_2$ (by Proposition 3 below) and $(2, d_2, d_3) \in \text{mdeg}(\text{Tame}(\mathbb{C}^3))$ for all $d_3 \geq d_2 \geq 2$ ([3] Corollary 2.3), then the assumption $d_3 \geq p_2 > p_1 \geq 3$ is not restrictive.

Key words and phrases. polynomial automorphism, tame automorphism, multidegree.
2000 Mathematics Subject Classification: 14Rxx, 14R10.

2. PROOF OF THE THEOREM

First, we recall one classical result (due to Sylvester) from the number theory, particularly from so-called coin problem or Frobenius problem [2].

Theorem 2. *If a, b are positive integers such that $\gcd(a, b) = 1$, then for every integer $k \geq (a-1)(b-1)$ there are $k_1, k_2 \in \mathbb{N}$ such that*

$$k = k_1 a + k_2 b.$$

Moreover $(a-1)(b-1) - 1 \notin a\mathbb{N} + b\mathbb{N}$.

In the proof we will, also, use the following proposition.

Proposition 3. ([3], Proposition 2.2) *If for a sequence of integers $1 \leq d_1 \leq \dots \leq d_n$ there is $i \in \{1, \dots, n\}$ such that*

$$d_i = \sum_{j=1}^{i-1} k_j d_j \quad \text{with } k_j \in \mathbb{N},$$

then there exists a tame automorphism F of $\mathbb{C}^n$ with $\text{mdeg } F = (d_1, \dots, d_n)$.

By the above proposition, in order to prove Theorem 1, it is enough to show that if $d_3 \notin p_1\mathbb{N} + p_2\mathbb{N}$, then $(p_1, p_2, d_3) \notin \text{mdeg}(\text{Tame}(\mathbb{C}^3))$.

In the proof of the above implication we will use some results and notions from the paper of Shestakov and Umirbayev [5, 6].

The first one is the following

Definition 1. ([5], Definition 1) *A pair $f, g \in k[X_1, \dots, X_n]$ is called $*$ -reduced if*
 (i) f, g are algebraically independent;
 (ii) $\bar{f}, \bar{g}$ are algebraically dependent, where $\bar{h}$ denotes the highest homogeneous part of h ;
 (iii) $\bar{f} \notin k[\bar{g}]$ and $\bar{g} \notin k[\bar{f}]$.

Definition 2. ([5], Definition 1) *Let $f, g \in k[X_1, \dots, X_n]$ be a $*$ -reduced pair with $\deg f < \deg g$. Put $p = \frac{\deg f}{\gcd(\deg f, \deg g)}$. In this situation the pair f, g is called p -reduced pair.*

Theorem 4. ([5], Theorem 2) *Let $f, g \in k[X_1, \dots, X_n]$ be a p -reduced pair, and let $G(x, y) \in k[x, y]$ with $\deg_y G(x, y) = pq + r, 0 \leq r < p$. Then*

$$\deg G(f, g) \geq q(p \deg g - \deg g - \deg f + \deg[f, g]) + r \deg g.$$

In the above theorem $[f, g]$ means the Poisson bracket of f and g , but for us it is only important that

$$\deg[f, g] = 2 + \max_{1 \leq i < j \leq n} \deg \left(\frac{\partial f}{\partial x_i} \frac{\partial g}{\partial x_j} - \frac{\partial f}{\partial x_j} \frac{\partial g}{\partial x_i} \right)$$

if f, g are algebraically independent, and $\deg[f, g] = 0$ if f, g are algebraically dependent.

Notice, also, that the estimation from Theorem 4 is true even if the condition (ii) of Definition 1 is not satisfied. Indeed, if $G(x, y) = \sum_{i,j} a_{i,j} x^i y^j$, then, by the algebraic independence of $\bar{f}$ and $\bar{g}$ we have:

$$\begin{aligned} \deg G(f, g) &= \max_{i,j} \deg(a_{i,j} f^i g^j) \geq \deg_y G(x, y) \cdot \deg g = \\ &= (qp + r) \deg g \geq q(p \deg g - \deg f - \deg g + \deg[f, g]) + r \deg g. \end{aligned}$$

The last inequality is a consequence of the fact that $\deg[f, g] \leq \deg f + \deg g$.

We will also use the following theorem.

Theorem 5. ([5], Theorem 3) *Let $F = (F_1, F_2, F_3)$ be a tame automorphism of $\mathbb{C}^3$. If $\deg F_1 + \deg F_2 + \deg F_3 > 3$ (in other words if F is not a linear automorphism), then F admits either an elementary reduction or a reduction of types I-IV (see [5] Definitions 2-4).*

Let us, also, recall that an automorphism $F = (F_1, F_2, F_3)$ admits an elementary reduction if there exists a polynomial $g \in \mathbb{C}[x, y]$ and a permutation σ of the set $\{1, 2, 3\}$ such that $\deg(F_{\sigma(1)} - g(F_{\sigma(2)}, F_{\sigma(3)})) < \deg F_{\sigma(1)}$.

Proof. (of Theorem 1) Assume that $F = (F_1, F_2, F_3)$ is an automorphism of $\mathbb{C}^3$ such that $\text{mdeg } F = (p_1, p_2, d_3)$. Assume, also, that $d_3 \notin p_1\mathbb{N} + p_2\mathbb{N}$. By Theorem 2 we have:

$$(1) \quad d_3 < (p_1 - 1)(p_2 - 1).$$

First of all we show that this hypothetical automorphism F does not admit reductions of type I-IV.

By the definitions of reductions of types I-IV (see [5] Definitions 2-4), if $F = (F_1, F_2, F_3)$ admits a reduction of these types, then $2 \mid \deg F_i$ for some $i \in \{1, 2, 3\}$. Thus if d_3 is odd, then F does not admit a reduction of types I-IV. Assume that $d_3 = 2n$ for some positive integers n .

If we assume that F admits a reduction of type I or II, then by the definition (see [5] Definition 2 and 3) we have $p_1 = sn$ or $p_2 = sn$ for some odd $s \geq 3$. Since $p_1, p_2 \leq d_3 = 2n < sn$, then we obtain a contradiction.

And, if we assume that F admits a reduction of type III or IV, then by the definition (see [5] Definition 4) we have:

$$n < p_1 \leq \frac{3}{2}n, \quad p_2 = 3n$$

or

$$p_1 = \frac{3}{2}n, \quad \frac{5}{2}n < p_2 \leq 3n.$$

Since $p_1, p_2 \leq d_3 = 2n < \frac{5}{2}n, 3n$, then we obtain a contradiction. Thus we have proved that our hypothetical automorphism F does not admit a reduction of types I-IV.

Now we will show that it, also, does not admit an elementary reduction.

Assume, by a contrary, that

$$(F_1, F_2, F_3 - g(F_1, F_2)),$$

where $g \in k[x, y]$, is an elementary reduction of (F_1, F_2, F_3) . Then we have $\deg g(F_1, F_2) = \deg F_3 = d_3$. But, by Theorem 4, we have

$$\deg g(F_1, F_2) \geq q(p_1 p_2 - p_1 - p_2 + \deg[F_1, F_2]) + r p_2,$$

where $\deg_y g(x, y) = q p_1 + r$ with $0 \leq r < p_1$. Since F_1, F_2 are algebraically independent, $\deg[F_1, F_2] \geq 2$ and then

$$p_1 p_2 - p_1 - p_2 + \deg[F_1, F_2] \geq p_1 p_2 - p_1 - p_2 + 2 > (p_1 - 1)(p_2 - 1).$$

This and (1) follows that $q = 0$, and that:

$$g(x, y) = \sum_{i=0}^{p_1-1} g_i(x) y^i.$$

Since $\text{lcm}(p_1, p_2) = p_1 p_2$, then the sets

$$p_1 \mathbb{N}, p_2 + p_1 \mathbb{N}, \dots, (p_1 - 1)p_2 + p_1 \mathbb{N}$$

are disjoint. This follows that:

$$\deg \left(\sum_{i=0}^{p_1-1} g_i(F_1) F_2^i \right) = \max_{i=0, \dots, p_1-1} (\deg F_1 \deg g_i + i \deg F_2).$$

Since, also,

$$d_3 \notin \bigcup_{r=0}^{p_1-1} (rp_2 + p_1 \mathbb{N})$$

(because $d_3 \notin p_1 \mathbb{N} + p_2 \mathbb{N}$), then it is easy to see that

$$\deg \left(\sum_{i=0}^{p_1-1} g_i(F_1) F_2^i \right) = d_3$$

is impossible.

Now, assume that

$$(F_1, F_2 - g(F_1, F_3), F_3),$$

where $g \in k[x, y]$, is an elementary reduction of (F_1, F_2, F_3) . Since $d_3 \notin p_1 \mathbb{N} + p_2 \mathbb{N}$, $p_1 \nmid d_3$ and $\gcd(p_1, d_3) = 1$. This means, by Theorem 4, that

$$\deg g(F_1, F_3) \geq q(p_1 d_3 - d_3 - p_1 + \deg[F_1, F_3]) + r d_3,$$

where $\deg_y g(x, y) = qp_1 + r$ with $0 \leq r < p_1$. Since $p_1 d_3 - d_3 - p_1 + \deg[F_1, F_3] \geq p_1 d_3 - 2d_3 \geq d_3 > p_2$ and since we want to have $\deg g(F_1, F_3) = p_2$, then $q = r = 0$. This means that $g(x, y) = g(x)$. But since $p_2 \notin p_1 \mathbb{N}$, then equality $\deg g(F_1) = p_2$ is impossible.

Finally, if we assume that $(F_1 - g(F_2, F_3), F_2, F_3)$ is an elementary reduction of (F_1, F_2, F_3) , then in the same way as in the previous case we obtain a contradiction. $\square$

3. SOME CONSEQUENCES

Theorem 6. *Let $p_2 > 3$ be prime number and $d_3 \geq p_2$ be integer. Then $(3, p_2, d_3) \in \text{mdeg}(\text{Tame}(\mathbb{C}^3))$ if and only if $d_3 \notin \{2p_2 - 3k \mid k = 1, \dots, \lfloor \frac{p_2}{3} \rfloor\}$.*

Proof. Since $p_2 > 3$ is a prime number, $p_2 \equiv 1 \pmod{3}$ or $p_2 \equiv 2 \pmod{3}$. Let $r \in \{1, 2\}$ be such that $p_2 \equiv r \pmod{3}$. It is easy to see that if $d_3 \geq p_2$ and $d_3 \equiv 0 \pmod{3}$ or $d_3 \equiv r \pmod{3}$, then $d_3 \in 3\mathbb{N} + p_2 \mathbb{N}$. Thus, by Theorem 2

$$2(p_2 - 1) - 1 \neq 0, r \pmod{3}.$$

Take any d_3 such that $p_2 \leq d_3 \leq 2p_2 - 3$ and $d_3 \not\equiv 0, r \pmod{3}$. Since $d_3 \leq 2p_2 - 3$ and $d_3 \equiv 2p_2 - 3 \pmod{3}$, then one can see that $d_3 \notin 3\mathbb{N} + p_2 \mathbb{N}$, because in other case we would have $2p_2 - 3 \in 3\mathbb{N} + p_2 \mathbb{N}$ which is a contradiction with Theorem 2. Thus

$$\begin{aligned} \{d_3 &\in \mathbb{N} \mid d_3 \geq p_2, d_3 \notin 3\mathbb{N} + p_2 \mathbb{N}\} = \\ &= \{d_3 \in \mathbb{N} \mid p_2 \leq d_3 \leq 2p_2 - 3, d_3 \equiv 2p_2 - 3 \pmod{3}\} \\ &= \{2p_2 - 3k \mid k = 1, \dots, \lfloor \frac{p_2}{3} \rfloor\} \end{aligned}$$

$\square$

Theorem 7. (a) If $d_3 \geq 7$, then $(5, 7, d_3) \notin \text{mdeg}(\text{Tame}(\mathbb{C}^3))$ if and only if $d_3 \neq 8, 9, 11, 13, 16, 18, 21, 23$.

(b) If $d_3 \geq 11$, then $(5, 11, d_3) \notin \text{mdeg}(\text{Tame}(\mathbb{C}^3))$ if and only if $d_3 \neq 12, 13, 14, 17, 18, 19, 23, 24, 28, 29, 34, 39$.

(c) If $d_3 \geq 13$, then $(5, 13, d_3) \notin \text{mdeg}(\text{Tame}(\mathbb{C}^3))$ if and only if $d_3 \neq 14, 16, 17, 19, 21, 22, 24, 27, 29, 32, 34, 37, 42, 47$.

(d) If $d_3 \geq 11$, then $(7, 11, d_3) \notin \text{mdeg}(\text{Tame}(\mathbb{C}^3))$ if and only if $d_3 \neq 12, 13, 15, 16, 17, 19, 20, 23, 24, 26, 27, 30, 31, 34, 37, 38, 41, 45, 48, 52, 59$.

Proof. This is a consequence of Theorem 2 and Theorem 1. For example to prove (a), by Theorem 2 and Theorem 1 we only have to check which of the numbers $7, 8, \dots, 23 = (5-1)(7-1) - 1$ are elements of the set $5\mathbb{N} + 7\mathbb{N}$. $\square$

REFERENCES

- [1] H.W.E. Jung, *Über ganze birationale Transformationen der Ebene*, J. reine angew. Math. 184 (1942), 161-174.
- [2] A. Brauer, *On a problem on partitions*, Amer. J. Math. 64 (1942), 299-312,
- [3] M. Karaś, *There is no tame automorphism of $\mathbb{C}^3$ with multidegree $(3, 4, 5)$* , to appear
- [4] W. van der Kulk, *On polynomialrings in two variables*, Nieuw Archief voor Wiskunde (3) 1 (1953), 33-41.
- [5] I.P. Shestakov, U.U. Umirbayev, *The Nagata automorphism is wild*, Proc. Natl. Acad. Sci. USA 100 (2003), 12561-12563.
- [6] I.P. Shestakov, U.U. Umirbayev, *The tame and the wild automorphisms of polynomial rings in three variables*, J. Amer. Math. Soc. 17 (2004), 197-227.

MAREK KARAŚ
 INSTYTUT MATEMATYKI
 UNIwersYTETU Jagiellońskiego
 UL. Łojasiewicza 6
 30-348 KRAKÓW
 POLAND
 e-mail: Marek.Karas@im.uj.edu.pl